

Melolonthidae (Coleoptera: Scarabaeoidea) of the Palaearctic and Oriental Regions.

II. Annotated catalogue of the genus *Sophrops* Fairmaire, 1887

Marek Bunalski 

Department of Entomology and Environment Protection, University of Life Sciences, Dąbrowskiego 159, 60-594 Poznań, Poland.

marek.bunalski@up.poznan.pl

Abstract: Based on the analysis of the literature data and specimens from museum and private collections the checklist of species belonging to the genus *Sophrops* Fairmaire, 1887 has been collated. Each species is presented with its full synonymy, bibliographical data, and distribution in alphabetical order. For 15 species, a new taxonomic status has been proposed. The new distributional data of some species are given.

Keywords: Rhizotroginae, chorology, bibliography, new status, new data

Introduction

The genus *Sophrops* was established by Fairmaire in 1887 for *Sophrops parviceps* from Yunnan. Its laconic description and imprecise genus characteristics caused difficulties in classifying subsequent species to this genus. This accounted for researchers including newly discovered species in better described genera, such as *Holotrichia* Hope, 1837 and *Brahmina* Blanchard, 1851, or created new genera, e.g. *Microtrichia* Brenske, 1900. A significant change was brought only with Arrow's work (1946), who by synonymising the genus *Microtrichia* transferred the species that were included in it to the genus *Sophrops*, thus establishing the morphological profile of the genus *Sophrops* through the descriptions of subsequent species.

The first checklist for the genus *Sophrops* was created by Frey (1972b), thereby giving a stimulus to further research (Frey 1973a,b, 1975a; Nomura 1977a,b). Efforts to systematise the genus *Sophrops* significantly accelerated in 1990s thanks to a group of scientists from different countries (Itoh 1992, 1993, 1994, 1995; Gu & Zhang 1992, 1995, 1996; Bunalski 1995) and they continue to this

day (Keith 2005, Matsumoto 2015, 2018, 2020).

Materials and methods

The following study is based on the analysis of the original descriptions, redescriptions, and taxonomic commentaries, as well as specimens from entomological collections. The species distribution was determined with the help of faunistic and faunistic-ecological studies, as well as lists and catalogues, which are cited next to particular species. All species are accompanied with bibliographical data, complete synonymy, as well as distribution in alphabetical order.

Furthermore the new distributional data of some species are given. The new data coming from collections:

DEEP – Department of Entomology and Environment Protection, University of Life Sciences, Poznań, Poland;

MIZ – Museum and Institute of Zoology, Polish Academy of Sciences, Warszawa, Poland;

MLU – Museum of Lund University, Lund, Sweden;

NHM – Natural History Museum, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, Poland;

ZMUK – Zoological Museum University of København, København, Danmark.

Results

The presented catalogue consists of 125 species from the genus *Sophrops*, including 15 species for which new taxonomic statuses have been proposed.

Genus *Sophrops* Fairmaire, 1887

= *Microtrichia* Brenske, 1900: 343.

= *Eretusa* Reitter, 1902: 174.

= *Noanda* Reitter, 1902: 174.

- ***Sophrops abscessus* (Brenske, 1892)**

Brahmina abscessa Brenske, 1892a: 102; Dalla Torre 1912: 220.

Microtrichia abscessa: Moser 1913b: 272.

Sophrops abscessa: Arrow 1946: 23; Frey 1972b: 354.

Distribution: LAOS. VIETNAM: Ho-Shi-Minh (= Saigon, loc. typ.).

- ***Sophrops acalcarius* Gu & Zhang, 1996**

Sophrops acalcarium Gu & Zhang, 1996: 23; Smetana & Král 2006: 227.

Sophrops acalcarius: Bezděk 2016: 278.

Distribution: CHINA: Guandong prov., Guangzhou (loc. typ.).

- ***Sophrops acutangularis* (Moser, 1918)**

Microtrichia acutangularis Moser, 1918: 243.

Sophrops acutangularis: Frey 1972b: 354; Smetana & Král 2006: 227; Bezděk 2016: 278.

Distribution: CHINA: Jiangxi prov., Pingshiang (loc. typ.).

- ***Sophrops analesetalis* Gu & Zhang, 1992**

Sophrops analesetalis Gu & Zhang, 1992: 257; Bezděk 2016: 278.

Distribution: CHINA: Yunnan prov. (loc. typ.).

- ***Sophrops andamanus* (Brenske, 1899) n. comb.**

Holotrichia andamana Brenske, 1899: 178; Dalla Torre 1912: 200; Frey 1971b: 222; Rao et al. 2013: 22.

Lachnosterna andamana: Barlow 1900: 257.

Microtrichia andamana: Brenske 1900a: 342; Moser 1913b: 272.

Distribution: INDIA: Andaman Arch. (loc. typ.).
Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops arrowi* Frey, 1973**

Sophrops arrowi Frey, 1973: 236.

Distribution: MYANMAR: Lashio (loc. typ.).

- ***Sophrops assamensis* (Moser, 1913) n. comb.**

Brahmina assamensis Moser, 1913a: 62; Sabatinelli 1992: 603.

Microtrichia assamensis: Moser 1913b: 272.

Distribution: INDIA: Assam prov., Cachar (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops basicollis* Arrow, 1946**

Sophrops basicollis Arrow, 1946: 24; Frey 1972b: 354.

Distribution: MYANMAR: Kambaiti (loc. typ.).

- ***Sophrops bifenestratus* Arrow, 1946**

Sophrops bifenestratus Arrow, 1946: 23.

Sophrops bifenestrato: Frey 1972b: 354.

Distribution: MYANMAR: Katha Mohnyin (loc. typ.), Sadon.

New localities: North THAILAND, Doi Inthanon N.P., Mt. Doi Inthanon, 2500 m, V 1998, 1 female, leg. W. Czajczyński (coll. DEEP).

- ***Sophrops bituberculatus* (Moser, 1908)**

Brahmina bituberculata Moser, 1908b: 359; Dalla Torre 1912: 221.

Microtrichia bituberculata: Moser 1913b: 272.

Sophrops bituberculatus: Arrow 1946: 23; Frey 1972b: 354; Itoh 1994: 300.

= *Sophrops kadleci* Frey, 1969: 110.

Distribution: LAOS (loc. typ.). THAILAND: Lampang prov.: Doi Pa Muang, Hang Chat; Nan prov.: Sanean Salai.

New data: N. THAILAND, Doi Inthanon N.P., Chiang Mai, V 1998, 2500 m, 1 female, leg. W. Czajczyński (coll. DEEP); N. THAILAND, Koh Chang, 4 IV 1959, 1 male, leg. B. Degerbol (coll. MLU).

• ***Sophrops brevisetalis* Gu & Zhang, 1992**

Sophrops brevisetalis Gu & Zhang, 1992: 258; Bezděk 2016: 278.

Distribution: CHINA: Hubei prov. (loc. typ.).

• ***Sophrops brunneus* (Moser, 1915)**

Microtrichia brunnea Moser, 1915b: 585.

Sophrops brunnea: Frey 1972b: 354; Smetana & Král 2006: 227.

Sophrops brunneus: Itoh 1994: 306; Bezděk 2016: 278.

Distribution: CHINA: Yunnan prov. (loc. typ.). LAOS: Samneua. THAILAND: Chiang Mai, Doi Chiang Khian, Doi Suthep, Erawan Resort, Nan vic., Phuping Palace, Sanean Salai.

• ***Sophrops burmanicus* (Moser, 1913)**

Brahmina burmanica Moser, 1913a: 67.

Microtrichia burmanica: Moser 1913b: 272.

Sophrops burmanica: Frey 1972b: 354; Sabatinelli 1992: 618; Smetana & Král 2006: 227.

Sophrops burmanicus: Bezděk 2016: 278.

Distribution: BHUTAN. MYANMAR: Schwegyin, Teinzo (loc. typ.), Tharrawaddy.

• ***Sophrops buruensis* (Brenske, 1892) n. comb.**

Brahmina buruensis Brenske, 1892a: 114; Dalla Torre 1912: 221.

Microtrichia buruensis: Moser 1913b: 272.

Distribution: INDONESIA: Maluku Arch., Buru Isl. (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

• ***Sophrops cantonensis* Petrovitz, 1969**

Sophrops cantonensis Petrovitz, 1969: 215; Bezděk 2016: 278.

Distribution: CHINA: Guangxi prov., Guangdong, Wuchow (loc. typ.).

• ***Sophrops cardoni* (Brenske, 1892)**

Brahmina Cardoni Brenske, 1892a: 108; Barlow 1900: 264; Dalla Torre 1912: 221.

Microtrichia Cardoni: Moser 1913b: 272.

Brahmina cardoni: Sabatinelli 1992: 603; Smetana & Král 2006: 212.

Sophrops cardoni: Bezděk 2016: 278.

Distribution: INDIA: Sikkim prov., Darjeeling distr., Kurseong (loc. typ.). NEPAL.

• ***Sophrops cephalotes* (Burmeister, 1855)**

Phytalus cephalotes Burmeister, 1855: 352; Gemminger and Harold 1869: 1172.

Holotrichia cephalotes: Brenske 1892a: 81; Moser 1908a: 341; Dalla Torre 1912: 201; Itoh 1995: 195.

Microtrichia cephalotes: Moser 1913b: 272; Zhang 1992: 501.

Sophrops cephalotes: Smetana & Král 2006: 227; Keith *et al.* 2012: 388; Bezděk 2016: 278; Rössner *et al.* 2017: 112; Matsumoto 2020: 352.

= *Holotrichia horishana*: Nijima and Kinoshita 1923: 222.

= *Holotrichia (Noanda) horishana*: Nomura 1977b: 90.

= *Holotrichia (Holotrichia) horishana*: Smetana & Král 2006: 218.

Distribution: CHINA: Guangxi prov.; Siuchuan prov., Hengduan Mts.; Yunnan prov., Taching Co., Mt. Gaoligangshan. GERMANY – introduced. LAOS: Samneua. TAIWAN: Bao Shan, Liukuei, Puli, Taipei, Wulai. THAILAND: Chum Phon, Fang, Khao Soi Dao. VIETNAM: Loel prov., Na Haeo, Chiang Mai, Chiem Hoa, Cuc Phuong, Mae Hong Son Mts., Mt. Mau Son, Tam Dao, Vinh Phu.

New data: TAIWAN, Soon Moon Lake, 3-12 VI 1995, 1 male (coll. DEEP).

- ***Sophrops chinensis* (Brenske, 1892)**

Brahmina chinensis Brenske, 1892a: 101; Moser 1908a: 342.

Sophrops chinensis: Reitter 1902: 173; Dalla Torre 1912: 249; Medvedev 1951: 273; Frey 1972b: 354; Bezděk 2016: 278.

Microtrichia chinensis: Moser 1913b: 272.

Distribution: CHINA (loc. typ.); VIETNAM: Mt. Mau Son.

- ***Sophrops confusus* (Moser, 1917)**

Microtrichia confusa Moser, 1917: 91.

Sophrops confusa: Frey 1972b: 354.

Distribution: INDONESIA: Sumatra Ins. (loc. typ.).

- ***Sophrops coronarius* Gu & Zhang, 1992**

Sophrops coronarius Gu & Zhang, 1992: 257; Bezděk 2016: 278.

Distribution: CHINA: Sichuan prov., Yunnan prov. (loc. typ.).

- ***Sophrops costatus* (Walker, 1859)**

Rhizotrogus costatus Walker, 1859: 54; Dalla Torre 1912: 228.

Microtrichia costata: Arrow 1916: 433.

Sophrops costatus: Bunalski 1995: 5.

= *Microtrichia puttalama*: Brenske 1900a: 360.

= *Holotrichia puttalama*: Dalla Torre 1912: 208.

= *Holotrichia puttalana* (sic!): Frey 1971b: 223.

Distribution: INDIA: Karnataka prov. SRI LANKA (= Ceylon, loc. typ.): Central Prov.: Nuwara-Eliya vic.; East Prov.: Inginiyagala; North-Central Prov.: Anuradhnapura vic.; Uva Prov.: Badulla vic., Bibile; West Prov.: Colombo vic.

- ***Sophrops cotesi* (Brenske, 1892)**

Brahmina Cotesi Brenske, 1892a: 100; Dalla Torre 1912: 221.

Lachnosterna cotesi: Barlow 1900: 259.

Sophrops cotesi: Arrow 1946: 23; Sabatinelli 1992: 618; Smetana & Král 2006: 227; Bezděk 2016: 279; Muraleedharan & Roy 2016: 364.

Holotrichia cotesi: Sabatinelli 1992: 609.

Sophrops cotesi (sic!): Thapa 2000: 206.

Distribution: BHUTAN. INDIA: Himachal Pradesh prov., Sikkim prov. (loc. typ.), Darjeeling distr. NEPAL: Kathmandu.

- ***Sophrops cribripennis* (Brenske, 1892)**

Brahmina cribripennis Brenske, 1892a: 106; Brenske 1894b: 279; Brenske 1900a: 147; Dalla Torre 1912: 221.

Microtrichia cribripennis: Moser 1913b: 272; Moser 1917: 91.

Sophrops cribripennis: Frey 1972b: 354; Itoh 1993: 267.

Distribution: INDONESIA: Sumatra Isl. (loc. typ.): Indrapura, Jija val., Soekaranda. MALAYSIA: Pehang st.: Cameron Highlands.

- ***Sophrops defreinae* Frey, 1973**

Sophrops defreinae Frey, 1973: 236; Sabatinelli 1992: 618; Smetana & Král 2006: 227; Bezděk 2016: 279.

Sophrops defreinae (sic!): Thapa 2000: 206.

Distribution: INDIA: Kumaon Himalaya distr.: Bhim val., Naini val. (loc. typ.); Uttaranchal prov.; Uttar Pradesh prov. NEPAL: Kathmandu.

- ***Sophrops deliensis* (Moser, 1917)**

Microtrichia deliensis Moser, 1917: 92.

Sophrops deliensis: Frey 1972b: 354; Itoh 1993: 267.

Distribution: INDONESIA: Sumatra Isl.: Deli (loc. typ.), Harau val., Bukit Tinggi vic., Lembah Anai, Paya Kumbuh.

New localities: INDONESIA, Sumatra, 18 VIII 1910, 2 male, 1 female (coll. DEEP).

- ***Sophrops densatus* Frey, 1971**

Sophrops densata Frey, 1971a: 124; Frey 1972b: 354.

Distribution: MYANMAR: Toungoo (loc. typ.).

- ***Sophrops dorsalis* (Frey, 1969) n. comb.**

Holotrichia dorsalis Frey, 1969: 112.

Distribution: VIETNAM: Dalat (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops egregius* (Moser, 1912)**

Holotrichia egregia Moser, 1912: 447.

Microtrichia egregia: Moser 1913b: 272.

Sophrops egregia: Frey 1972b: 354; Sabatinelli 1992: 618.

Distribution: INDIA: Assam prov. (loc. typ.).

- ***Sophrops eurystomus* (Burmeister, 1855)**

Phytalus eurystomus Burmeister, 1855: 253; Gemminger and Harold 1869: 1172.

Holotrichia eurystoma: Brenske 1892: 176; Frey 1971b: 223.

Microtrichia eurystoma: Brenske 1900a: 361; Dalla Torre 1912: 208; Arrow 1916: 433.

Sophrops eurystomus: Bunalski 1995: 5.

Rhizotrogus exactus: Walker 1859: 55.
= *Holotrichia parva*: Brenske 1892: 180.

Distribution: SRI LANKA (= Ceylon, loc. typ.): Sabaragamuwa Prov.: Rakwana vic.; South Prov.: Galle vic.; West Prov.: Colombo vic.

- ***Sophrops excisus* Itoh, 1994**

Sophrops excisus Itoh, 1994: 309.

Distribution: THAILAND: Chiang Khan, Chiang Mai, Doi Pui, Doi Suthep (loc. typ.).

New data: North THAILAND, Doi Suthep, 19-20 V 1958, 1 male, 3 female, leg. B. Degerbol (coll. MLU).

- ***Sophrops flabellatus* (Brenske, 1892) n. comb.**

Brahmina flabellata Brenske, 1892a: 104; Dalla Torre 1912: 222; Sabatinelli 1992: 604.

Microtrichia flabellata: Moser 1913b: 272.

Distribution: INDIA: Sikkim prov. (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops formosanus* (Moser, 1913)**

Microtrichia formosana Moser, 1913b: 273; Nijjima & Kinoshita 1923: 54.

Sophrops formosana: Frey 1972b: 354; Nomura 1977b: 94; Smetana & Král 2006: 227.

Sophrops formosanus: Bezděk 2016: 279.

Distribution: TAIWAN: Fenchihu, Hotso, Jiuyuehtan, Lienhwachi, Kenting, Kuantzungling,

Nanshanchi, Shitoshan, Taipei, Tainan (loc. typ.), Taitung, Wushe.

New data: TAIWAN (Formosa), A Li Shan, 17-26 VI 1995, 1 male (coll. DEEP).

- ***Sophrops foveatus* (Moser, 1909)**

Brahmina foveata Moser, 1909: 476; Dalla Torre 1912: 222.

Microtrichia foveata: Moser 1913b: 272.

Sophrops foveata: Frey 1972b: 354.

Sophrops foveatus: Itoh 1994: 304.

Distribution: LAOS: Samneua. THAILAND: Kanchanaburi prov.: Koh Anem vill.; Lampang prov.: Doi Pa Muang; Lang Phrae prov.: Maeta Teak Rep. Forest; Chiang Mai vic., Doi Chiang Khian, Fang, Saiyok vic. VIETNAM (= Tonkin, loc. typ.).

New data: NW VIETNAM, Sông Hồng riv.val., Lao Cai, 5 V 1960, 1m, leg. K. Galewski; idem, 16-17 V 1960, 1 male, 1 female, leg. K. Galewski; idem, 7-8 VI 1960, 1 female, leg. K. Galewski - at the foot of the busches hills, at light. (coll. MIZ), idem, 12 IV 1961, 2 male, 1 female, leg. R. Bartke (coll. NHM); North THAILAND, Doi Suthep, 27 IV 1958, 1 male, 1 female, leg. B. Degerbol, idem, 19-23 V 1958, 3 male, 3 female, leg. B. Degerbol; North THAILAND, Doi Inthanon, 15 V 1958, 2 male, leg. B. Degerbol; Fang, 24-25 V 1958, 2 male, 1 female, leg. B. Degerbol (coll. MLU); THAILAND, Doi Inthanon N.P., Chiang Mai prov., Mae Klang, 10 X 1981, 1 female (coll. ZMUK).

- ***Sophrops furcifer* (Arrow, 1916)**

Microtrichia furcifer Arrow, 1916: 433.

Sophrops furcifer: Frey 1972b: 354.

Distribution: SRI LANKA (= Ceylon, loc. typ.).

- ***Sophrops hauseri* Balthasar, 1932**

Sophrops hauseri Balthasar, 1932: 20; Frey 1972b: 354; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov.: Soling-ho riv. val. (loc. typ.).

- ***Sophrops heteropygus* (Moser, 1913)**

Microtrichia heteropyga Moser, 1913a: 57.

Sophrops heteropygus: Arrow 1946: 21.

Sophrops heteropyga: Sabatinelli 1992: 618; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: INDIA: Assam prov.: Manipur; Sikkim prov.: Darjeeling distr. (loc. typ.); West Bengal prov. MYANMAR: Sadon.

- ***Sophrops heydeni* (Brenske, 1892)**

Brahmina Heydeni Brenske, 1892a: 108.

Sophrops Heydeni: Reitter 1902: 173; Dalla Torre 1912: 249; Nijima & Kinoshita 1927: 25.

Sophrops heydeni: Medvedev 1951: 272; Murayama 1954: 101; Stebnicka 1980: 264; Smetana & Král 2006: 227; Kim 2011: 142; Bezděk 2016: 279; Bezborodov & Zinchenko 2018: 14.

Microtrichia Heydeni: Moser 1913b: 272.

Sophrops hegdeni (sic!): Frey 1972b: 354.

Distribution: CHINA: Fujian prov., Guangdong prov., Heilongjiang prov., Liaoning prov., Shanxi prov. NORTH KOREA (loc. typ.): Chungcheong buk-do prov., Chungcheong nam-do prov., Ganwon-do prov., Gyeosangbuk-do, Gyeonggi-do prov., Gyeosang nam-do prov., Jeju-do prov., Hamgyeong buk-do prov., Hamgyeong nam-do prov., Hwanghae buk-do prov., Jeolla buk-do prov., Jeolla nam-do prov., Jeju-do prov., Pyeongan buk-do prov., Pyeongan nam-do prov. RUSSIA: Khabarovskii Krai, Primorskii Krai. SOUTH KOREA.

New data: RUSSIA, Primorskii Krai, Lesozavodskii Raion, Filaretovka, 6-10 VIII 1992, 1 male, leg. A. Jadwiszczak (coll. DEEP).

- ***Sophrops himalayicus* (Brenske, 1892)**

Brahmina himalayica Brenske, 1892a: 113; Dalla Torre 1912: 222.

Microtrichia himalayica: Moser 1913b: 272.

Sophrops himalayica: Frey 1972b: 354; Sabatinelli 1992: 618; Smetana & Král 2006: 227.

Sophrops himalayicus: Bezděk 2016: 279.

Distribution: INDIA: Sikkim prov.: Darjeeling distr. (loc. typ.).

- ***Sophrops impressicollis* (Moser, 1913)**

Microtrichia impressicollis Moser, 1913b: 274.

Sophrops impressicollis: Frey 1972b: 354; Itoh 1993: 266.

Distribution: INDONESIA: Java Isl.: West Preanger, Gunong Tji Salimar; Sumatra Isl.: Bedagei (loc. typ.), Deli.

- ***Sophrops iridescens* (Moser, 1908)**

Holotrichia iridescens Moser, 1908a: 340; Dalla Torre 1912: 203; Chang 1965: 40; Smetana & Král 2006: 220; Bezděk 2016: 268.

Sophrops iridescens: Keith *et al.* 2012: 388.

Distribution: CHINA: Yunnan prov.: Mt. Gaoligongshan. VIETNAM: Mt. Mau Son, Than-Moi (loc. typ.).

- ***Sophrops iridipennis* (Brenske, 1892)**

Holotrichia iridipennis Brenske, 1892b: 177; Dalla Torre 1912: 203; Frey 1971b: 221-222; Sabatinelli 1992: 610; Smetana & Král 2006: 227; Bezděk 2016: 268.

Lchnosterna iridipennis: Barlow 1900: 261.

Microtrichia iridipennis: Moser 1913b: 272.

Sophrops iridipennis: Muraleedharan & Roy 2016: 364.

Distribution: INDIA: Seebis (loc. typ.), Assam st., Karnataka st., Kerala st., Maharashtra st., Meghalaya st., Khasi Hills, Orissa st., Tamil Nadu st. MYANMAR.

- ***Sophrops irregularis* Arrow, 1946**

Sophrops irregularis Arrow, 1946: 22; Frey 1972b: 354.

Distribution: MYANMAR: Kambaiti (loc. typ.).

- ***Sophrops kanarana* (Moser, 1918)**

Microtrichia kanarana Moser, 1918: 243.

Sophrops kanarana: Frey 1972b: 354.

Distribution: INDIA: Karnataka st.: Kanara Arch. (loc. typ.).

- ***Sophrops karschi* (Brenske, 1892)**

Holotrichia Karschi Brenske, 1892b: 179; Dalla Torre 1912: 203.

Microtrichia Karschi: Brenske 1900a: 342; Moser 1913b: 272.

Sophrops karschi: Frey 1972b: 354; Gite *et al.* 2015: 1271-1272; Bajad *et al.* 2017: 1365.

Holotrichia karschi: Matsumoto 2014: 56.

Distribution: INDIA: Pondicherry (loc. typ.), Karnataka st., Kerala st., Maharashtra st., Tamil Nadu st.

- ***Sophrops kaszabi* (Frey, 1972) n. comb.**

Holotrichia kaszabi Frey, 1972a: 117.

Distribution: INDIA: Kerala st.: Calicut distr. (loc. typ.).

Note: Morphological characters correspond to the genus *Sophrops*.

- ***Sophrops kawadai kawadai* (Nomura, 1959)**

Microtrichia kawadai Nomura, 1959: 48.

Sophrops kawadai: Nomura 1964: 54; Nomura 1965: 3; Frey 1972b: 354.

Sophrops kawadai kawadai: Nomura 1977a: 77; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: JAPAN: Ryukyu Arch.: Amami-Oshima Isl. (loc. typ.), Iriomote Isl.: Mt. Hateruma.

- ***Sophrops kawadai okinawaensis* Nomura, 1977**

Sophrops kawadai okinawaensis Nomura, 1977a: 77; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: JAPAN: Ryukyu Arch.: Okinawa-shoto Isl., Shuri (loc. typ.).

- ***Sophrops keralensis* Frey, 1973**

Sophrops keralensis Frey, 1973: 238.

Distribution: INDIA: Kerala st., Trivandram DC Poonmudi Range (loc. typ.).

- ***Sophrops konishii konishii* Nomura, 1970**

Sophrops konishii Nomura, 1970: 67.

Sophrops konishii konishii: Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: JAPAN: Ryukyu Arch.: Iriomote Isl., Mt. Hateruma (loc. typ.), Ishigaki Isl., Mt. Omoto.

- ***Sophrops konishii yonaguniensis* Nomura, 1970**

Sophrops konishii subsp. yonaguniensis Nomura, 1970: 68.

Sophrops konishii yonaguniensis: Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: JAPAN: Yonaguni-shima Isl., Sonai (loc. typ.).

- ***Sophrops latericostatus* (Fairmaire, 1889) n. comb.**

Phytalus? latericostatus Fairmaire, 1889: 340.

Brahmina latericostata: Brenske 1892a: 115; Dalla Torre 1912: 222.

Distribution: VIETNAM (= Tonkin, loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops laticeps* (Moser, 1913)**

Microtrichia laticeps Moser, 1913b: 275.

Sophrops laticeps: Frey 1972b: 355.

Distribution: INDONESIA: Java Isl. (loc. typ.).

- ***Sophrops latiusculus* Nomura, 1977**

Sophrops latiuscula Nomura, 1977b: 93.

Sophrops latiuscula: Smetana & Král 2006: 227.

Sophrops latiusculus: Bezděk 2016: 279.

Distribution: TAIWAN: Kukwan (loc. typ.).

- ***Sophrops latus* Frey, 1972**

Sophrops lata Frey, 1972a: 113; Frey 1972b: 354; Smetana & Král 2006: 227.

Sophrops latus: Bezděk 2016: 279.

Distribution: CHINA: Fujian prov.: Kuantan (loc. typ.).

- ***Sophrops longicarinatus* (Brenske, 1892)**

Holotrichia longicarinata Brenske, 1892b: 181; Dalla Torre 1912: 204.

Lachnosterna longicarinata: Barlow 1900: 259.

Microtrichia longicarinata: Moser 1913b: 272.
Sophrops longicarinatus: Arrow 1946: 23.
 Distribution: INDONESIA: Sumatra Isl.: Kallies (loc. typ.).

- ***Sophrops longiflabella* Gu & Zhang 1995**

Sophrops longiflabella Gu & Zhang, 1995: 9;
 Smetana & Král 2006: 227; Bezděk 2016: 279.
 Distribution: CHINA: Hubei prov. (loc. typ.).

- ***Sophrops luangia* (Moser, 1912) n. comb.**

Holotrichia luangia Moser, 1912: 448.
Microtrichia luangia: Moser 1913b: 272.
 Distribution: VIETNAM: Luang Tri (loc. typ.).
 Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops luridipennis* (Moser, 1915)**

Microtrichia luridipennis Moser, 1915a: 134.
Sophrops luridipennis: Frey 1972b: 355;
 Smetana & Král 2006: 227; Bezděk 2016: 279.
 Distribution: CHINA: Yunnan prov. (loc. typ.).

- ***Sophrops microphyllus* (Brenske, 1892)**

Brahmina microphylla Brenske, 1892a: 104;
 Dalla Torre 1912: 222.
Microtrichia microphylla: Moser 1913b: 272.
Sophrops microphylla: Frey 1972b: 355.
 Distribution: INDIA: Karnataka st.: Belgaum (loc. typ.). THAILAND: Bangkok.

- ***Sophrops mindanaoensis* Moser, 1924**

Microtrichia mindanaoensis Moser, 1924: 356.
Sophrops mindanaoensis: Frey 1972b: 355.
 Distribution: PHILIPPINES: Mindanao Isl. (loc. typ.).

- ***Sophrops monticola* (Moser, 1915) n. comb.**

Holotrichia monticola Moser, 1915b: 583.
 Distribution: PHILIPPINES: Luzon Isl.: Mt. Makiling (loc. typ.).
 Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops montivagus* (Moser, 1915)**

Microtrichia montivaga Moser, 1915b: 587.
Sophrops montivaga: Arrow 1946: 22; Frey 1972b: 355.
 Distribution: VIETNAM: Mt. Mau Son (loc. typ.).

- ***Sophrops nigrofuscus* (Moser, 1921)**

Microtrichia nigrofusca Moser, 1921: 66.
Sophrops nigrofusca: Frey 1972b: 355.
 Distribution: MYANMAR: Rangun (loc. typ.).

- ***Sophrops nilgirianus* (Moser, 1913)**

Microtrichia nilgiriana Moser, 1913a: 59.
Sophrops nilgiriana: Frey 1972b: 355.
 Distribution: INDIA: Tamil Nadu st.: Nilgiri Hills (loc. typ.).

- ***Sophrops nubiliventris* (Bates, 1891) n. comb.**

Lachnosterna nubliliventris Bates, 1891a: 16.
Holotrichia nubliliventris: Dalla Torre 1912: 204; Frey 1971b: 221; Sabatinelli 1992: 611; Smetana & Král 2006: 220; Bezděk 2016: 268.
Microtrichia nubliliventris: Moser 1913b: 272.
 Distribution: INDIA: Himachal Pradesh st.: Kulu (loc. typ.).
 New data: NE INDIA, Darjiling prov., Galatea, 1 female; Himalaya Mts., 1 male (coll. DEEP).
 Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops nudus* (Moser, 1915)**

Brahmina nuda Moser, 1915a: 140; Smetana & Král 2006: 214; Bezděk 2016: 257.
Sophrops nuda: Frey 1972b: 355.
 Distribution: CHINA: Fujian prov. (loc. typ.).

- ***Sophrops obscurus* (Brenske, 1892) n. comb.**

Holotrichia obscura Brenske, 1892b: 162; Dalla Torre 1912: 223; Sabatinelli 1992: 611.
Lachnosterna obscura: Barlow 1900: 260.
Microtrichia obscura: Moser 1913b: 272.
 Distribution: INDIA: Assam st.: Sikkim prov. (loc. typ.). NEPAL.

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops opacidorsalis* Itoh, 1994**

Sophrops opacidorsalis Itoh, 1994: 308.

Distribution: THAILAND: Chiang Mai vic., Doi Mae Salong, Doi Pui (loc. typ.); Doi Chiang Khian, Doi Ang Khang, Surat Thani, Khao Poata, Nan prov., Na Noi, Mt. Doi Ku Sathan.

- ***Sophrops pallidus* Itoh, 1994**

Sophrops pallidus Itoh, 1994: 311.

Distribution: THAILAND: Fang (loc. typ.), Nan prov., Khao Yai, Wiang Sa.

- ***Sophrops parviceps* Fairmaire, 1887**

Sophrops parviceps Fairmaire, 1887: 106; Brenske 1892a: 116; Reitter 1902: 173; Dalla Torre 1912: 249; Frey 1972b: 355; Keith *et al.* 2012: 388; Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov. (loc. typ.).

- ***Sophrops paucisetosa* Frey, 1971**

Sophrops paucisetosa Frey, 1971a: 126; Frey 1972b: 355.

Distribution: LAOS: Vientiane (loc. typ.). THAILAND: Chiang Mai. VIETNAM: Dilinh.

- ***Sophrops peronosporus* Gu & Zhang, 1992**

Sophrops peronosporus Gu & Zhang, 1992: 256; Bezděk 2016: 279.

Distribution: CHINA: Guangdong prov., Hunan prov. (loc. typ.), Sichuan prov.

- ***Sophrops pexicollis* (Fairmaire, 1886)**

Schizonycha (*Holotrichia*) *pexicollis* Fairmaire, 1886: 324.

Lachnosterna pexicollis: Bates 1891b: 75.

Holotrichia pexicollis: Brenske 1898: 347; Dalla Torre 1912: 205; Smetana & Král 2006: 220.

Sophrops pexicollis: Keith 2005: 98; Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov. (loc. typ.).

- ***Sophrops planicollis* (Burmeister, 1855)**

Phytalus planicollis Burmeister, 1855: 352; Gemminger & Harold 1869: 1172.

Holotrichia planicollis: Moser 1908a: 341; Dalla Torre 1912: 205; Itoh 1995: 195.

Micritrichia planicollis: Moser 1913b: 272.

Sophrops planicollis: Frey 1971b: 206; Smetana & Král 2006: 227; Bezděk 2016: 279; Matsumoto 2020: 54.

= *Ancylonycha nigra*: Redtenbacher 1867: 67.

= *Holotrichia* (*Eretusa*) *nigra*: Reitter 1902: 174; Dalla Torre 1912: 204; Medvedev 1951: 295; Smetana & Král 2006: 218.

Distribution: CHINA: Hongkong (loc. typ.), Tschushan, Jianxi prov., Jian, Sichuan prov., Zhejiang prov. LAOS: Xieng Khouang vic. THAILAND: Chiang Mai, Doi Chiang Khian, Doi Pui, Doi Suthep. VIETNAM: Babe, Hanoi vic., Mt. Mau Son, Chiem Hoa, Sapa.

- ***Sophrops planipennis* (Moser, 1913)**

Microtrichia planipennis Moser, 1913b: 272.

Sophrops planipennis: Frey 1972b: 355.

Distribution: INDONESIA: Borneo Isl., Kalimantan prov., Mahakam (loc. typ.).

- ***Sophrops problematicus* (Brenske, 1899) n. comb.**

Holotrichia problematica Brenske, 1899: 178; Dalla Torre 1912: 205; Frey 1971b: 221; Sabatinelli 1992: 611; Bezděk 2016: 269.

Lachnosterna problematica: Barlow 1900: 260.

Microtrichia problematica: Brenske 1900a: 342; Moser 1913b: 272.

Distribution: INDIA: Kashmir reg., Shrinagar (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

- ***Sophrops promeanus* (Moser, 1918)**

Microtrichia promeana Moser, 1918: 244.

Sophrops promeana: Frey 1972b: 355.

Sophrops promeanus: Itoh 1994: 307.

Distribution: MYANMAR: Prome (loc. typ.). THAILAND: Kanchanaburi, Pu Nam Long Hot Spring. VIETNAM.

New data: North THAILAND, Doi Inthanon N.P., Chiang Mai, V 1998, 1 male, leg. W. Czajczyński (coll. DEEP).

- ***Sophrops pruinosipennis* (Moser, 1915)**

Microtrichia pruinosipennis Moser, 1915b: 586.

Sophrops pruinosipennis: Frey 1972b: 355.

Distribution: LAOS (loc. typ.).

- ***Sophrops pruinosipyga* Gu & Zhang, 1996**

Sophrops pruinosipyga Gu & Zhang, 1996: 25; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: CHINA: Anhui prov.: Huangshan (loc. typ.), Sichuan prov.: Baoxing.

- ***Sophrops pseudoplanicollis* Matsumoto, 2020**

Sophrops pseudoplanicollis Matsumoto, 2020: 52.

Distribution: THAILAND: Kamphaeng Phet (loc. typ.), Klong Wang Jao, Samoeng Prov., Chiang Mai.

- ***Sophrops pubiventris* (Burmeister, 1855)**

Phytalus pubiventris Burmeister, 1855: 353.

Brahmina pubiventris: Brenske 1892a: 109; Dalla Torre 1912: 223.

Microtrichia pubiventris: Moser 1913b: 272.

Sophrops pubiventris: Frey 1972b: 355.

Distribution: INDONESIA: Java Isl. (loc. typ.).

- ***Sophrops pumilus* (Sharp, 1881)**

Lachnosterna pumila Sharp, 1881: 225.

Brahmina pumila: Brenske 1892a: 110; Brenske 1894b: 279; Moser 1908a: 343; Dalla Torre 1912: 205.

Holotrichia pumila: Brenske 1900b: 148; Dalla Torre 1912: 223.

Microtrichia pumila: Moser 1913b: 272.

Sophrops pumilus: Arrow 1946: 22.

Sophrops pumila: Frey 1972b: 355.

= *Phyllophaga quedana* Saylor, 1937: 320.

Distribution: INDONESIA: Java Isl., Sumatra Isl.: Rawas distr., Soeroelengoen (loc. typ.), Indrapura, Silago, Soekaranda. VIETNAM: Mt. Mau Son.

- ***Sophrops purkynei* Balthasar, 1932**

Sophrops Purkyněi Balthasar, 1932: 21.

Sophrops purkunei: Frey 1972b: 355; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov., Soling-ho riv. val. (loc. typ.).

- ***Sophrops reticulata* Frey, 1969**

Sophrops reticulata Frey, 1969: 109; Frey 1972b: 355.

Distribution: LAOS: Fanky, Vientiane vic. (loc. typ.).

- ***Sophrops roeri* Frey, 1972**

Sophrops roeri Frey, 1972a: 112; Frey 1972b: 355; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: CHINA: Fujian prov.: Kuatun, Shaowu (loc. typ.).

- ***Sophrops rotundicollis* Itoh, 1994**

Sophrops rotundicollis Itoh, 1994: 300.

Distribution: THAILAND: Chiang Mai, Doi Pui, Doi Sang (loc. typ.), Fang.

- ***Sophrops rufofulvus* (Moser, 1919)**

Microtrichia rufoflava Moser, 1919: 31.

Sophrops rufofulva: Frey 1972b: 355; Sabatinelli 1992: 618.

Distribution: INDIA: Assam st., Meghalaya st.: Khasi Hills (loc. typ.).

- ***Sophrops rugatus* (Moser, 1915)**

Microtrichia rugata Moser, 1915b: 588.

Sophrops rugata: Frey 1972b: 355; Smetana & Král 2006: 227.

Sophrops rugatus: Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov. (loc. typ.).

- ***Sophrops rugifrons* (Moser, 1913)**

Microtrichia rugifrons Moser, 1913a: 58.

Sophrops rugifrons: Frey 1972b: 355.

Distribution: INDIA: Belganne (= Belgaum, loc. typ.).

- ***Sophrops rugipennis* (Frey, 1972)**

Metabolus rugipennis Frey, 1972a: 111.

Sophrops rugipennis: Gu & Zhang 1992: 257; Smetana & Král 2006: 277; Bezděk 2016: 279.

Distribution: CHINA: Fujian prov., Kuantun (loc. typ.).

- ***Sophrops rugulosus confusus* (Moser, 1917)**

Microtrichia confusa Moser, 1917: 91.

Sophrops confusa: Frey 1972b: 354.

Sophrops rugulosa confusa: Itoh 1993: 262.

Distribution: INDONESIA: Sumatra Isl. (loc. typ.): Barisan Mts., Bukit Tinggi vic., Air Mancur, Mt. Singgalang, Paya Kumbuh, Pangkalan.

- ***Sophrops rugulosus harauensis* Itoh, 1993**

Sophrops rugulosa harauensis Itoh, 1993: 264.

Distribution: INDONESIA: Sumatra Isl., Paya Kumbuh, Harau val. (loc. typ.).

- ***Sophrops rugulosus rugulosus* (Brenske, 1892)**

Brahmina rugulosa Brenske, 1892a: 107; Dalla Torre 1912: 223.

Microtrichia rugulosa: Moser 1913b: 272.

Sophrops rugulosa: Frey 1972b: 355.

Sophrops rugulosa rugulosa: Itoh 1993: 262.

Distribution: INDONESIA: Borneo Isl., Balaikrangan, Mt. Bawang. MALAYSIA: Borneo Isl., Sarawak prov. (loc. typ.).

- ***Sophrops schereri* Frey, 1975**

Sophrops schereri Frey, 1975a: 258.

Distribution: INDONESIA: Sumatra Isl. (loc. typ.).

- ***Sophrops sculpticollis* (Blanchard, 1850)**

Ancylonycha sculpticollis Blanchard, 1851: 139.

Lachnosterna sculpticollis: Gemminger and Harold 1869: 1169.

Holotrichia sculpticollis: Dalla Torre 1912: 206; Frey 1971b: 221-222.

Sophrops sculpticollis: Bajad et al. 2017: 1365.

= *Microtrichia densicollis* Moser, 1913a: 56.

Distribution: INDIA: Bombay (loc. typ.), Karnataka st., Kerala st., Maharashtra st., Tamil Nadu st., Nilgiri Hills.

New data: INDIA, Madras st., Shevaroy hills, Yeseand Galen, 1 female, Koefod Missionar (coll. DEEP); INDIA, Tamil Nadu st., Nilghiri hills, Kotagiri 15 km SE., Kunjappanai env., 900 m, 22-30 V 1999, 1 female, leg. Z. Kejval & M. Tryzna (coll. DEEP).

- ***Sophrops semiserrata* (Brenske, 1894)**

Holotrichia semiserrata Brenske, 1894a: 70; Dalla Torre 1912: 206; Smetana & Král 2006: 220; Bezděk 2016: 269.

Microtrichia semiserrata: Moser 1913b: 272.

Sophrops semiserrata: Matsumoto 2018: 351.

Distribution: CHINA: Shanghai (loc. typ.).

- ***Sophrops semitomentosus* (Frey, 1975)**

Holotrichia semitomentosa Frey, 1975b: 317.

Sophrops semitomentosus: Bunalski 1985: 6.

Distribution: SRI LANKA: Bolawatta (loc. typ.).

- ***Sophrops sericeicollis* (Moser, 1915)**

Microtrichia sericeicollis Moser, 1915b: 585.

Sophrops sericeicollis: Arrow 1946: 21.

Sophrops sericeicollis (sic!): Frey 1972b: 355.

Distribution: MYANMAR: Myitkyina vic. VIETNAM: Mt. Mau Son (loc. typ.).

- ***Sophrops setosus* (Brenske, 1892)**

Brahmina setosa Brenske, 1892a: 99; Dalla Torre 1912: 223.

Microtrichia setosa: Moser 1913b: 272.

Sophrops setosa: Frey 1972b: 355; Sabatinelli 1992: 618; Smetana & Král 2006: 227.

Sophrops setosus: Bezděk 2016: 279.

= *Phyllophaga quetosis* Saylor, 1937: 320.

Distribution: BHUTAN. INDIA: Sikkim prov., Darjeeling distr. (loc. typ.).

- ***Sophrops sharpi* (Brenske, 1892)**

Holotrichia Sharpi Brenske, 1892b: 180; Dalla Torre 1912: 206.

Microtrichia Sharpi: Brenske 1900a: 342; Moser 1913b: 272.

Sophrops sharpi: Matsumoto 2020: 55.

= *Holotrichia deplanata* Brenske, 1900b: 150.

Distribution: INDONESIA: Borneo Isl., Sarawak st. (loc. typ.); Sumatra Isl., Batoe Bara, Bedagei, Indrapura. MALAYSIA: Borneo Isl., Penang (loc. typ.), Malacca, Perak, Selangor Ulu. SINGAPORE.

• ***Sophrops shillongensis* (Brenske, 1899) n. comb.**

Brahmina shillongensis Brenske, 1899: 177; Dalla Torre 1912: 223; Sabatinelli 1992: 605.

Microtrichia shillongensis: Moser 1913b: 272.

Distribution: INDIA: Meghalaya prov.: Shillong (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

• ***Sophrops siamensis* (Brenske, 1892)**

Brahmina siamensis Brenske, 1892a: 103; Barlow 1900: 264; Dalla Torre 1912: 223; Sabatinelli 1992: 605.

Microtrichia siamensis: Moser 1913b: 272.

Sophrops siamensis: Frey 1972b: 355.

Distribution: MALAYSIA: Malacca. THAILAND (= Siam, loc. typ.).

• ***Sophrops signatifrons* (Moser, 1913)**

Microtrichia signatifrons Moser, 1913a: 57.

Sophrops signatifrons: Frey 1972b: 355; Sabatinelli 1992: 619.

Distribution: INDIA: Assam st., Lakhimpur Hills, Dajoo (loc. typ.).

• ***Sophrops sikkimensis* (Brenske, 1892) n. comb.**

Brahmina sikkimensis Brenske, 1892a: 113; Dalla Torre 1912: 223; Sabatinelli 1992: 606; Smetana & Král 2006: 213; Bezděk 2016: 257.

Microtrichia sikkimensis: Moser 1913b: 272.

Distribution: INDIA: Sikkim prov. (loc. typ.), Darjeeling distr.

Note: morphological characters correspond to the genus *Sophrops*.

• ***Sophrops similis* Arrow, 1946**

Sophrops similis Arrow, 1946: 21; Frey 1972b: 355.

Distribution: MYANMAR: Kambaiti (loc. typ.).

• ***Sophrops simplex* Frey, 1971**

Sophrops simplex Frey, 1971a: 125; Frey 1972b: 355.

Brahmina simplex: Smetana & Král 2006: 213; Bezděk 2016: 257.

Distribution: LAOS: Vientiane (loc. typ.).

VIETNAM: Dilinh (loc. typ.).

• ***Sophrops singhalensis* (Brenske, 1892)**

Holotrichia singhalensis Brenske, 1892b: 178; Brenske 1899: 179; Frey 1971b: 222.

Microtrichia singhalensis: Brenske 1900b: 360; Dalla Torre 1912: 208; Arrow 1916: 433.

Sophrops singhalensis: Bunalski 1995: 5.

Distribution: SRI LANKA (= Ceylon, loc. typ.): Sabaragamuwa prov., Rakwana; West prov., Colombo.

• ***Sophrops sjoestedti* (Moser, 1921)**

Microtrichia Sjöstedt: Moser, 1921: 68.

Sophrops sjoestedti: Frey 1972b: 355.

Distribution: LAOS. THAILAND (= Siam, loc. typ.).

• ***Sophrops standfussi* (Brenske, 1892)**

Holotrichia Standfussi Brenske, 1892b: 182; Brenske 1894b: 279; Dalla Torre 1912: 206.

Lachnosterna Standfussi: Barlow 1900: 262.

Holotrichia (Noanda) Standfussi: Reitter 1902: 174.

Microtrichia Standfussi: Moser 1913b: 272.

Sophrops standfussi: Matsumoto 2015: 9.

Distribution: INDONESIA: Sumatra Isl. (loc. typ.): Aceh vic., Bandar Baru, Bukit Tinggi vic., Harau Valley, Paya Kumbuh, Soekaranda.

• ***Sophrops stenocarpus* Gu & Zhang, 1996**

Sophrops stenocarpus Gu & Zhang, 1996: 24; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: CHINA: Sichuan prov., Jiulong (loc. typ.).

- ***Sophrops striatus* (Brenske, 1892)**

Brahmina striata Brenske, 1892a: 112.

Microtrichia striata: Moser 1913b: 272.

Sophrops striata: Reitter 1902: 173; Dalla Torre 1912: 249; Nijijima & Kinoshita 1927: 24; Medvedev 1951: 274; Murayama 1954: 99; Frey 1972b: 355; Stebnicka 1980: 265; Smetana & Král 2006: 227; Kim 2011: 143.

Sophrops striatus: Bezděk 2016: 279.

Distribution: CHINA (loc. typ.): Henan prov. NORTH KOREA: Chungcheong nam-do prov., Gangwon-do prov., Gyeonggi-do prov., Gyeongsang nam-do prov., Hvanghe nam-do prov., Jeolla nam-do prov, Jeju-do prov., Kangwon-do prov., Pyeong-an nam-do prov., Phjongan nam-do prov. SOUTH KOREA.

New data: North KOREA, Mjohjang-san Mts., 17 VI 1983, 1 male; idem, 21 VI 1985, 1 male, Exped. ISEZ Cr.; North KOREA, Mjohjang-san Mts., Hjangsan-chón riv., 20-25 VI 1981, 1 male, leg. A. Szeptycki (coll. NHM).

- ***Sophrops subrugatus* (Moser, 1921)**

Microtrichia subrugata Moser, 1921: 67.

Sophrops subgrata (sic!): Frey 1972b: 355.

Sophrops subrugata: Smetana & Král 2006: 227.

Sophrops subrugatus: Bezděk 2016: 279.

Distribution: CHINA: Guandong prov., Guandong (= Canton, loc. typ.).

- ***Sophrops subrugipennis* (Moser, 1921)**

Microtrichia subrugipennis Moser, 1921: 67.

Sophrops subrugipennis: Frey 1972b: 355.

Distribution: INDONESIA: Borneo Isl., Doesonlanden (loc. typ.).

- ***Sophrops subrugosus* (Moser, 1915)**

Microtrichia subrugosa Moser, 1915a: 133.

Sophrops subrugosa: Frey 1972b: 356; Smetana & Král 2006: 227.

Sophrops subrugosus: Bezděk 2016: 279.

Distribution: CHINA: Kiukiang (loc. typ.), Jiangxi prov.

- ***Sophrops sumatrensis* (Brenske, 1892)**

Brahmina sumatrensis Brenske, 1892a: 105; Dalla Torre 1912: 223.

Microtrichia sumatrensis: Moser 1913b: 272.

Sophrops sumatrensis: Frey 1972b: 356.

= *Phyllophaga sumatricollis* Saylor, 1937: 320.

Distribution: INDONESIA: Sumatra Isl. (loc. typ.).

- ***Sophrops taiwanus* Nomura, 1977**

Sophrops taiwana Nomura, 1977b: 93; Smetana & Král 2006: 227.

Sophrops taiwanus: Bezděk 2016: 279.

Distribution: TAIWAN: Liehwachi (loc. typ.), Nantou Hsien, Jenai, Wushe.

- ***Sophrops takatoshii* Itoh, 1990**

Sophrops takatoshii Itoh, 1990: 9; Smetana & Král 2006: 227; Bezděk 2016: 279.

Distribution: JAPAN: Ryukyu Arch., Miyakojima Isl. (loc. typ.).

- ***Sophrops tavoyensis* (Brenske, 1892)**

Brahmina tavoyensis Brenske, 1892a: 104; Barlow 1900: 265; Dalla Torre 1912: 223.

Microtrichia tavoyensis: Moser 1913b: 272.

Sophrops tavoyensis: Arrow 1946: 22; Frey 1972b: 356.

Distribution: MYANMAR: Mali Kyun Isl. (= Tavoy, loc. typ.).

- ***Sophrops teinzoanus* (Moser, 1918)**

Microtrichia teinzoana Moser, 1918: 245.

Sophrops teinzoana: Frey 1972b: 356.

Distribution: MYANMAR: Teinzo (loc. typ.).

- ***Sophrops thoracicus* (Brenske, 1892)**

Brahmina thoracica Brenske, 1892a: 109; Brenske 1896: 151; Barlow 1900: 265; Dalla Torre 1912: 223.

Microtrichia thoracica: Moser 1913b: 272.

Sophrops thoracica: Frey 1972b: 356; Sabatinelli 1992: 619; Smetana & Král 2006: 227.

Sophrops thoracicus: Bezděk 2016: 279.

= *Phyllophaga thoracalis* Saylor, 1937: 320.

Distribution: INDIA: Meghalaya st., Khasi Hills (loc. typ.); Sikkim prov., Darjeeling distr., Kurseong.

• ***Sophrops tonkinensis* (Moser, 1908)**

Brahmina tonkinensis Moser, 1908a: 342; Dalla Torre 1912: 223.

Microtrichia tonkinensis: Moser 1913b: 272.

Sophrops tonkinensis: Frey 1972b: 356; Itoh 1994: 310.

= *Phyllophaga tonkinana* Saylor, 1937: 320.

Distribution: THAILAND: Chiang Mai vic, Doi Chiang Khian, Doi Sang, Khao Yai. VIETNAM: Mt. Mau Son (loc. typ.), Vinh Phu, Tam Dao.

• ***Sophrops umbilicatus* Frey, 1973**

Sophrops umbilicatus Frey, 1973: 235; Bezděk 2016: 279.

Sophrops umbilicata: Frey 1975c: 241; Sabatinelli 1992: 619; Smetana & Král 2006: 227.

Distribution: BHUTAN. INDIA: Uttarkhand st., Kumaon Himalaya distr., Bhim val., Naini val. (loc. typ.).

• ***Sophrops unguicularis* (Moser, 1912) n. comb.**

Holotrichia unguiculari: Moser, 1912: 449.

Microtrichia unguicularis: Moser 1913b: 272.

Distribution: BRUNEI (loc. typ.).

Note: morphological characters correspond to the genus *Sophrops*.

• ***Sophrops vethi* (Moser, 1914)**

Microtrichia Vethi Moser, 1914: 152.

Sophrops Vethi: Frey 1972b: 356.

Distribution: VIETNAM: Hanoi (loc. typ.).

• ***Sophrops yangbiensis* Gu & Zhang, 1996**

Sophrops yangbiensis Gu & Zhang, 1996: 26; Bezděk 2016: 279.

Distribution: CHINA: Yunnan prov., Yangbi (loc. typ.).

References

- Arrow G. 1916. The Melolonthine Beetles of Ceylon. *The Annals and Magazin of Natural History*, ser. 8, 18: 429–444.
- Arrow G. 1946. Entomological results from the Swedish Expedition 1934 to Burma and British India. Coleoptera: Melolonthinae, collected by Rene Malaise. *Arkiv för Zoologi*, 38A (9): 1–33.
- Bajad VV, Dadmal SM, Undirwade DB. 2017. Diversity of white grubs (Coleoptera: Scarabaeidae: Melolonthinae) from Maharashtra. *Agriculture Update*, 12: 1363–1366.
- Balthasar V. 1932. Zwei neue Arten der Gattung *Sophrops* Fairm. aus Sud-China. (13. Beitrag zur Kenntnis der Scarabaeidae des palaearktischen Faunengebietes). *Wiener entomologische Zeitung*, 49: 19–22.
- Barlow E. 1900. List of the Melolonthini contained in the collection of the Indian Museum. *Indian Museum Notes*, 4 (5): 234–273.
- Bates HW. 1891a. Coleoptera from Kulu in N.W. India. *The Entomologist*, 24 (Suppl.): 7–22.
- Bates HW. 1891b. Coleoptera collected by Mr. Pratt on Upper Yang-Tsze, and on the borders of Tibet. Second Notice. Journey of 1890. *The Entomologist*, 24 (Suppl.): 69–80.
- Bezborodov VG, Zinchenko VK. 2018. New data on distribution of *Sophrops heydeni* (Brenske, 1892) (Coleoptera: Scarabaeidae) in Russia. *Far East Entomologist*, 352: 14–15.
- Bezděk A. 2016. Tribe Rhizotrogini Burmeister, 1855. In: Löbl I, Löbl D. (Eds). *Catalogue of Palaearctic Coleoptera. Vol. 3. Scarabaeoidea – Scirtoidea – Dascilloidea – Buprestoidea – Byrrhoidea. Revised and Updated Edition*. Leiden – Boston, pp. 249–280.
- Blanchard CE. 1851. (new taxa). In: Milne-Edwards H. (Ed.). *Catalogue de la Collection Entomologique du Muséum d'Histoire Naturelle de Paris. Classe des Insectes*.

- Ordre des Coléoptères. Tome 2.* Gide et Baudry, Paris, pp. 129–240.
- Brenske E. 1892a. Die Arten der Coleopteren-Gattung *Brahmina* Bl. *Berliner entomologische Zeitschrift*, 37 (1): 79–124.
- Brenske E. 1892b. Neue Arten der Coleopteren-Gattung *Holotrichia* (Lachnosterna). *Berliner entomologische Zeitschrift*, 37 (2): 159–192.
- Brenske E. 1894a. Die Melolonthiden der palaearktischen und orientalischen Region im Königlichen naturhistorischen Museum zu Brüssel. Beschreibung neuer Arten und Bemerkungen zu bekannten. *Mémoires de la Société Entomologique de Belgique*, 2: 3–87.
- Brenske E. 1894b. Zur Kenntnis der Melolonthiden Sumatra's. *Stettiner entomologische Zeitung*, 55: 274–280.
- Brenske E. 1896. Insectes du Bengale (37^e Mémoire.) Melolonthidae. *Annales de la Société Entomologique de Belgique*, 40: 150–164.
- Brenske E. 1898. Melolonthiden aus Africa. *Stettiner entomologische Zeitung*, 59: 333–394.
- Brenske E. 1899. Diagnoses Melolonthidarum novarum ex India orientali. *Indian Museum Notes*, 4: 176–179.
- Brenske E. 1900a. Contribution a l'étude de Faune de Sumatra (Cote Ouest - Vice-residence de Painan). VIII. Melolonthiden gesammelt von J-L Weyers. *Mémoires de la Société Entomologique de Belgique*, 7: 141–155.
- Brenske E. 1900b. Die Melolonthiden Ceylon's, unter besondere Berücksichtigung der von Dr. Horn gesammelten Arten. *Stettiner entomologische Zeitung*, 61: 341–361.
- Bunalski M. 1995. Melolonthinae of Sri Lanka (Coleoptera: Scarabaeoidea: Melolonthidae) with notes on the systematic position of some taxa. *Stobaeana*, 4: 1–6.
- Burmeister HCC. 1855. *Handbuch der Entomologie. Vierter Band. Besondere Entomologie. Fortsetzung. Zweite Abtheilung. Coleoptera Lamellicornia Phyllophaga chaenochela.* Theod. Chr. Friedr. Enslin, Berlin, IV, 2, I-X + 569 pp.
- Chang YW. 1965. Revision of Chinese may beetles of the genus *Holotrichia* Hope. III. (Coleoptera: Scarabaeidae). *Acta Zootaxonomica Sinica*, 2 (1): 37–56. [in Chinese]
- Dalla Torre KW. von 1912. Scarabaeidae: Melolonthinae III. In: Junk W, Schenkling S. (Eds). *Coleopterorum Catalogus auspiciis et auxillio W. Junk editus a S. Schenkling.* Berlin, 20, 49: 135–290.
- Fairmaire L. 1886. Descriptions de Coleopteres de l'intérieur de la Chine. *Annales de la Société Entomologique de France*, 6 (6): 303–356.
- Fairmaire L. 1887 Coléoptères de l'intérieur de la Chine (3e partie). *Annales de la Société Entomologique de Belgique*, 31: 87–136.
- Fairmaire L. 1889 (1888). Descriptions de coléoptères de l'Indo-Chine. *Annales de la Société Entomologique de France*, 6(8): 333–378.
- Frey G. 1969. Neue lamellicornia aus Südostasien und Australien (Col.). *Entomologische Arbeiten aus dem Museum G. Frey*, 20: 106–116.
- Frey G. 1971a. Neue Ruteliden und Melolonthiden aus Indien und Indochina (Col.). *Entomologische Arbeiten aus dem Museum Georg Frey*, 22: 109–133.
- Frey G. 1971b. Bestimmungstabelle der indischen und ceylonischen Arten der Gattung *Holotrichia* Hope (Col., Melolonth.). *Entomologische Arbeiten aus dem Museum G. Frey*, 22: 206–225.
- Frey G. 1972a. Neue chinesische Melolonthiden aus Museum Koenig in Bonn und einige neue *Holotrichia*-Arten (Col., Scarab., Melolonthinae). *Entomologische Arbeiten aus dem Museum G. Frey*, 23: 108–121.
- Frey G. 1972b. Katalog aller in der Gattung *Sophrops* Fairm. eingereichten Arten (Col. Scarab.). *Entomologische Arbeiten aus dem Museum G. Frey*, 23: 354–356.

- Frey G. 1973. Neue orientalische Melolonthiden (Col., Scarab.). *Entomologische Arbeiten aus dem Museum Georg Frey*, 24: 232–239.
- Frey G. 1975a. Neue Melolonthiden und Ruteliden aus den Beständen der Zoologischen Staatssammlung München. *Entomologische Arbeiten aus dem Museum G. Frey*, 26: 256–262.
- Frey G. 1975b. Zwei neue Holotrichia-Arten aus Ceylon und Südindien. *Entomologische Arbeiten aus dem Museum G. Frey*, 26: 316–318.
- Gemminger M, Harold B. von 1869. *Catalogus Coleopterorum, hucusque descriptorum, synonymicus et systematicus. Tom. IV. Scarabaeidae*. E.H. Gummi, Monachii: 979–1346 pp.+7 pp.
- Gite RB., Mohite PB., Rathour B. 2015. Distribution and diversity of white grub species in Western Maharashtra, India. *International Journal of Science and Research*, 4 (12): 1270–1272.
- Gu Y, Zhang YW. 1992. Description of four new species in the genus *Sophrops* Fairmaire (Scarabaeidae Coleoptera). *Journal of Shenyang Agricultural University*, 23: 256–259.
- Gu Y, Zhang YW. 1995. Revision of two genera of Melolonthidae with description of three new species. *Entomological Journal of East China*, 4 (2): 4–10.
- Gu Y, Zhang YW. 1996. Four new species of the genus *Sophrops* (Coleoptera: Melolonthidae) from China. *Entomotaxonomia*, 18: 23–31.
- Hope FW. 1837. *The Coleopterist's Manual, Containing the Lamellicorn Insects of Linneus and Fabricius*. Henry G. Bohn, London, 121 pp.+4 pp.
- Itoh T. 1990. A new species on the genus *Sophrops* (Melolonthinae, Scarabaeidae) from Is. Miyakolima, Ryukyus. *Lamellicornia*, 5: 9–10.
- Itoh T. 1993. A taxonomical note on the Sumatran species of the genus *Sophrops* (Melolonthidae, Melolonthinae, Melolonthini). *Elytra*, 21 (2): 259–269.
- Itoh T. 1994. Melolonthidae (Coleoptera) from Thailand, I. *Elytra*, 22 (2): 299–313.
- Itoh T. 1995. Melolonthidae (Coleoptera) from Thailand, II. *Elytra*, 23 (2): 195–204.
- Keith D. 2005. Sur quelques Scarabaeoidea paléarctiques et orientaux. *Bulletin Mensuel de la Société Linnéenne de Lyon*, 74 (3): 93–102.
- Keith D, Li J, Zhang X, Huang L. 2012. Scarabaeoidea Pleurosticti récoltés sur le Mt. Gaoligongshan en République Populaire de Chine (Insecta: Coleoptera). *Veröffentlichungen des Naturkundemuseums Erfurt*, 31: 387–397.
- Kim JI. 2011. Insect Fauna of Korea. Arthropoda: Insecta: Coleoptera: Scarabaeoidea (Pleurosticti). *The Flora and Fauna of Korea*, 143, 18: 1–253.
- Matsumoto T. 2014. Redescription of the genus *Holotrichia* Hope (Coleoptera, Scarabaeidae, Melolonthinae). *Kogane*, 20: 53–57.
- Matsumoto T. 2015. Changes of the systematic positions of eight species and two subgenera of the genus *Holotrichia*. *Kogane*, 17: 5–10.
- Matsumoto T. 2018. Transfer of the taxonomic positions of some *Holotrichia* species. *Elytra, New Series*, 8 (2): 349–353.
- Matsumoto T. 2020. A new species and a new synonym of the genus *Sophrops*. *Kogane*, 23: 52–56.
- Medvedev SI. 1951. Zhestkokrylye. Platinchatousye (Scarabaeidae). Podsem. Melolonthinae, tch. 1 (khrushchi). *Fauna SSSR*, 10, 1: 1–512.
- Moser J. 1908a. Verzeichnis der von H. Fruhstorfer in Tonkin gesammelten Melolonthiden. *Annales de la Société Entomologique de Belgique*, 52: 325–343.
- Moser J. 1908b. Neue Melolonthiden und Cetoniden. *Annales de la Société Entomologique de Belgique*, 52: 353–367.
- Moser J. 1909. Neue Arten der Melolonthiden-Gattungen *Holotrichia* und *Brahmina*.

- Annales de la Société Entomologique de Belgique*, 53: 468–478.
- Moser J. 1912. Neue Arten der Melolonthiden-Gattungen *Holotrichia* und *Pentelia*. *Annales de la Société Entomologique de Belgique*, 56: 420–449.
- Moser J. 1913a. Neue indische Melolonthiden. *Deutsche entomologische Zeitschrift*: 49–70.
- Moser J. 1913b. Beitrag zur Kenntnis der Melolonthiden. (Col.). *Deutsche entomologische Zeitschrift*: 271–297.
- Moser J. 1914. Neue Melolonthiden-Arten aus der Sammlung des Herrn Dr. H. J. Veth (Haag). *Notes from the Leyden Museum*, 36: 148–156.
- Moser J. 1915a. Beiträge zur Kenntnis der Melolonthiden (Col.) IV. *Deutsche Entomologische Zeitschrift*: 113–151.
- Moser J. 1915b. Neue Melolonthiden und Cetoniden. *Deutsche entomologische Zeitschrift*: 579–605.
- Moser J. 1917. Beitrag zur Kenntnis der Melolonthiden (Col.) VI. *Stettiner entomologische Zeitung*, 78: 54–97.
- Moser J. 1918. Neue Melolonthiden aus der Sammlung des deutschen Entomologischen Museums zu Berlin-Dahlem (Col.). *Stettiner entomologische Zeitung*, 79: 209–247.
- Moser J. 1919. Beitrag zur Kenntnis der Melolonthiden. VII. (Fortsetzung.). *Deutsche entomologische Zeitschrift*: 1–34.
- Moser J. 1921. Beitrag zur Kenntniss der Melolonthiden (Col.) XII. *Stettiner entomologische Zeitung*, 82: 48–73.
- Moser J. 1924. New species of Melolonthid beetles from the Philippine Islands. *The Philippine Journal of Science*, 25: 349–357.
- Muraleedharan N, Roy S. 2016. Arthropod pests and natural enemy communities in tea ecosystems of India. In: Chakravarthy AK, Sridhara S. (Eds). *Economic and ecological significance of Arthropods in diversified ecosystems. Sustaining regulatory mechanisms*. Springer Science et Business Media, Singapore, pp. 361–392.
- Murayama J. 1954. *Icones Melolonthidarum Manchuriae et Koreae I*. Nihon Gakjuitsu Sinokai, 163 pp., 6 pls. [in Japanese]
- Nijijima Y, Kinoshita E. 1923. Die Untersuchungen über japonische Melolonthiden (II). (Melolonthiden Japans und ihre Verbreitung). *Research Bulletins of the College Experiment Forest, College of Agriculture Hokkaido Imperial University Sapporo*, 2 (2): 1–253. [in Japanese]
- Nijijima Y, Kinoshita E. 1927. Die Untersuchungen über japonische Melolonthiden (III). *Research Bulletins of the College Experiment Forest, College of Agriculture Hokkaido Imperial University Sapporo*, 4: 1–97. [in Japanese]
- Nomura S. 1959. Notes on the Japanese Scarabaeoidea (Coleoptera) with two subspecies from Formosa. *Tôhō-Gakuhô*, 9: 39–54.
- Nomura S. 1964. Some new forms of Scarabaeoidea from Loochoo Islands. *Entomological Review of Japan*, 16 (2): 50–60.
- Nomura S. 1965. Some new forms of Scarabaeoidea from Loochoo Islands. III. *Entomological Review of Japan*, 18 (1): 1–9.
- Nomura S. 1970. Notes on some Scarabaeid-beetles from Loochoo and Formosa. *Entomological Review of Japan*, 22 (2): 61–72.
- Nomura S. 1977a. Notes on some Scarabaeid-beetles from Japan and Taiwan. *Tôhō-Gakuhô*, 27: 77–84.
- Nomura S. 1977b. On the Melolonthini of Taiwan (Coleoptera, Scarabaeidae). *Tôhō-Gakuhô*, 27: 85–109.
- Petrovitz R. 1969. Zwei neue Melolonthinae aus der orientalischen Region (Coleoptera, Scarabaeidae). *Mitteilungen aus der zoologischen Sammlung des Museums für Naturkunde in Berlin*, 45 (2): 215–219.
- Rao DV, Chandra K, Devi K. 2013. *Endemic animals of Andaman and Nicobar Islands*. Government of India, Kolkata, 182 pp.
- Redtenbacher L. (1867) 1868. *Reise der Österreichischen Fregatte Novara um die*

- Erde in den Jahren 1857, 1858, 1859 unter der Befehlen des Commodore B. von Wüllerstorff-Urbair. Zoologie II Part. Coleopteren.* Carl Gerold's Sohn, Wien, 249 pp.
- Reitter E. (1901) 1902. Bestimmungs-Tabelle der Melolonthidae aus der europäischen Fauna und den angrenzenden Ländern, enthaltend die Gruppen der Pachydemini, Sericini und Melolonthini. Heft 50. *Verhandlungen des Naturforschenden Vereins in Brünn*, 40: 93–303.
- Rössner E, Keith D, Weigel A. 2017. Der Fund eines ostasiatischen Blatthornkäfers in Bayern (Coleoptera: Scarabaeidae, Rhizotrogini). *Entomologische Zeitschrift*, 127 (2): 112–114.
- Sabatinelli G. 1992. Catalogue of the Scarabaeoidea Melolonthidae from Himalaya and North India: Macroductylini, Hopliini, Diplotaxini, Melolonthini and Sericini (Coleoptera). *Memorie della Società Entomologica Italiana*, 71 (11): 593–636.
- Saylor LW. 1937. Necessary changes in status of important rhizotrogid genera (Col. Scarabaeidae). *Revista de Entomologia Rio de Janeiro*, 7 (2-3): 318–322.
- Sharp D. 1881. Descriptions of new species of Melolonthini and Rutelini, collected in the island of Sumatra during the Scientific Sumatra-expedition. *Notes Leyden Museum*, 3: 219-242.
- Smetana A, Král D. 2006. Tribe Rhizotrogini Burmeister, 1855. In: Löbl I, Smetana A. (Eds). *Catalogue of Palaearctic Coleoptera. Vol. III.* Apollo Books, Stenstrup, pp. 207-228.
- Stebnicka Z. 1980. Scarabaeoides of the Democratic People's Republic of Korea. *Acta Zoologica Cracoviensia*, 24 (5): 191–297.
- Thapa VK. 2000. *An inventory of Nepal's Insects, Vol. III.* IUCN Nepal, Kathmandu, I-XI + 475 pp.
- Walker F. 1859. Characters of some apparently undescribed Ceylon Insects. *The Annals and Magazine of Natural History*, ser. 3, 3: 50–56.
- Zhang YW. 1992. Coleoptera: Melolonthidae, Dynastidae. In: Chen S. (Ed.). *Insects of the Hengduan Mountains Region. Vol. 1.* Science Press, Beijing, pp. 499-503.

Received: 4.04.2022

Accepted: 13.07.2022

Published online: 30.09.2022